

Formatted: Font: 9 pt  
Formatted: Font: 9 pt

# CHAPTER 2. CLIMATE CHANGE AND ALASKA

The Arctic Climate Impact Assessment, a comprehensive assessment conducted by hundreds of scientists and indigenous peoples, projects increases in temperature, glacial melt, permafrost temperature, ocean acidification, and continued loss of sea ice extent and thickness in Arctic regions (ACIA 2004, 2005).

This chapter is not a product of the Adaptation Advisory Group. The information in this chapter provides a scientific backdrop for the recommendations and options for adaptation in each sector. It presents findings from the scientific community regarding observed and projected climatic changes for Alaska, and attendant potential impacts on the Alaskan economy and lifestyles.


Deleted: predicts  
Deleted: ongoing increases in concentrations of greenhouse gases are expected to lead to concomitant

Over the past 50 years, Alaska has warmed at more than twice the rate of the rest of the United States (Karl et al. 2009). As a result, climate change impacts are much more pronounced than in other regions. Higher temperatures are already contributing to earlier spring snowmelt, reduced sea ice, widespread glacier retreat, and permafrost warming. These observed changes are consistent with climate model projections of greater warming over Alaska, especially in winter, as compared with the rest of the country (Karl et al. 2009). Warmer temperatures are expected to be accompanied by overall drier conditions and reduced soil moisture, by changes in sea level, and other changes.

Changes in climate will pose challenges for Alaskans and the Alaska economy. Climate change will also provide economic opportunities due to a longer summer tourism season, increased navigation potential, and other changes. Addressing these challenges and taking advantage of the opportunities requires

understanding the likely changes in climate that Alaska is to experience, and the concomitant changes in natural resources, infrastructure, and other human and natural systems on which urban and rural communities depend. This chapter summarizes relevant literature on projected climate for Alaska and potential impacts on natural and human systems.

Figure 2-1. Total Change in Mean Annual Temperature (°F), 1949-2008



Deleted: likely  
Deleted:

## Climate Change Science and Projections for Alaska

Climate has historically played a key role in shaping Alaska's natural environment, a vast and varied land spanning 586,400 square miles and over 33,000 miles of coastline. Current and projected changes in climate, however, are occurring rapidly with impacts to society, the environment, infrastructure, and economic drivers throughout the state. While natural variability spread over many centuries is well-

Source: ACRC 2009.

Formatted: Font: 9 pt  
Formatted: Font: 9 pt  
Field Code Changed  
Formatted: Font: 9 pt

Formatted: Font: 9 pt  
Formatted: Font: 9 pt

1 documented, the present rate of change and global mean surface temperature are higher than during any  
 2 similar period since 1600 AD (Overpeck, et al. 1997). Alaskan climate is particularly sensitive to these  
 3 changes. Temperature and precipitation in Alaska are driven in large part by the state's high latitude and  
 4 corresponding seasonal extremes in solar radiation, the influence of ocean waters and sea ice, and variations  
 5 in elevation (ACRC 2009). A general understanding of projected conditions is critical to planning for the future  
 6 and anticipating the types of adaptation measures that should be developed.

7 **Temperature**

8 Despite the considerable interannual and interdecadal influence of large-scale regime shifts , most notably the  
 9 Pacific Decadal Oscillation (PDO), average annual temperatures in Alaska have risen by 3.1° F in the past 60  
 10 years (ACRC 2009). This is consistent with the general warming trend observed throughout the circumpolar  
 11 Arctic. The distribution across Alaska is displayed in Figure 2.1.

12 An analysis of mean seasonal and annual temperature data for Alaskan communities over the past 60 years  
 13 shows the greatest changes occurring during the winter months as depicted in Figure 2.2. Multiple  
 14 communities have experienced increases of greater than 8° F in the winter months. This poses significant  
 15 implications to the natural ecosystems and the built environment in Alaska.

16 **Precipitation and Hydrology**

17 Atmospheric temperature is positively  
 18 correlated with the amount of precipitable  
 19 water (PW) in the atmosphere, and thus  
 20 the potential for precipitation. During  
 21 1979 to 2005, precipitable water in the  
 22 atmosphere at the northern latitudes  
 23 increased by approximately 5 to 10% per  
 24 decade during spring, increased 5% per  
 25 decade in summer, and declined (over  
 26 land masses) in autumn and winter  
 27 (White et al. 2007). Available  
 28 precipitation gauge data from across the  
 29 northern latitudes (between 55° and  
 30 85°N) suggests precipitation has  
 31 increased during winter and spring, with  
 32 a slight decrease during autumn (e.g.,  
 33 Serreze et al. 2000).

34 Snow distribution and redistribution data  
 35 is both difficult to gather and the stations  
 36 that gather such data provide only  
 37 sparse coverage across Alaska.

38 Increases in winter precipitation could  
 39 lead to increased snowpack (Serreze et  
 40 al. 2000), however, winter melting events  
 41 and a shortening of the period of snow accumulation could have the opposite effect. If the latter conditions

Figure 2-2. Total Change in Mean Seasonal and Annual Temperature (°F), 1949 -2008

Total Change in Mean Seasonal and Annual Temperature (°F), 1949 - 2008						
Region	Location	Winter	Spring	Summer	Autumn	Annual
Arctic	Barrow	6.5	4.4	2.8	3.4	4.3
	Interior	Bettles	8.5	4.6	1.8	1.1
	Big Delta	9.2	3.5	1.2	-0.2	3.4
	Fairbanks	7.7	3.8	2.3	-0.4	3.3
	McGrath	7.4	4.8	2.7	0.6	3.9
West Coast	Kotzebue	6.6	1.8	2.5	1.6	3.1
	Nome	4.4	3.6	2.5	0.6	2.8
	Bethel	6.6	5.0	2.3	0.1	3.6
	King Salmon	8.1	4.7	1.8	0.6	3.8
	Cold Bay	1.5	1.8	1.8	0.9	1.5
	St Paul	1.0	2.4	2.8	1.3	1.9
Southcentral	Anchorage	6.8	3.6	1.6	1.4	3.1
	Talkeetna	8.9	5.4	3.1	2.4	5.0
	Gulkana	8.1	2.4	0.9	0	2.8
	Homer	6.3	4.0	3.4	1.7	3.9
	Kodiak	0.9	2.3	1.2	-0.4	1.0
Southeast	Yakutat	4.9	3.1	1.8	0.3	2.6
	Juneau	6.6	3.1	2.1	1.4	3.3
	Annette	3.9	2.5	1.7	0.2	2.1
Average		6.0	3.5	2.1	0.9	3.1

Alaska Climate Research Center Geophysical Institute, University of Alaska Fairbanks

Source: ACRC 2009

Formatted: Font: 9 pt  
Formatted: Font: 9 pt  
Field Code Changed  
Formatted: Font: 9 pt

Formatted: Font: 9 pt  
Formatted: Font: 9 pt

1 dominated and overall snowpack decreased, Alaska could expect a shorter spring melting period with lower  
2 runoff intensity, and generally lower summer baseflows. In either event, changes in timing, duration, thickness,  
3 and distribution of seasonal snow cover significantly impacts many aspects of the hydrologic cycle including  
4 surface runoff, groundwater recharge, and river streamflow (White et al. 2007).

5 In addition, rain-on-snow events can have a profound impact on wildlife. For example, when a heavy crust  
6 forms on snow after a rain event, caribou may be unable to access food and large-scale die-offs may occur.  
7 To complicate potential impacts of snowpack and river dynamics, glacial runoff has a significant impact on  
8 many Alaskan rivers. For example, Hinzman et al. (2005) found increasing trends in discharge from glacially  
9 fed Alaskan rivers and decreasing trends in nonglacially fed rivers. With continued warming, the glacial  
10 contribution to rivers could increase. This affect would then decrease with diminishing glaciers.

Deleted:  
Deleted:

11 The potential change in precipitation, snowpack, runoff, and riverflow could have many different effects on  
12 Alaskans. Due to the appreciable interannual variability and variation introduced by decadal cycles (i.e.,  
13 Pacific Decadal Oscillation), longer term trends in precipitation will be difficult to discern and prepare for.  
14 However, both decadal variability and longer term trends should be incorporated into long term planning.  
15 While most attention is paid to net annual or seasonal precipitation, the nature of the precipitation events is  
16 important to human and natural ecosystems. For example, an increase in storminess could have a dramatic  
17 impact on erosion and consequently infrastructure. Likewise, midwinter melting events could reduce the  
18 potential for spring flooding, particularly that flooding caused by ice jams on rivers. As with all climate impacts,  
19 changes will be regional in nature.

20 Warmer spring and fall weather will result in longer growing seasons and a higher number of growing degree  
21 days statewide. This in turn will increase water loss from soils through evapotranspiration. This drying effect  
22 will generally result in a net loss of moisture, despite increased rainfall.

23 In the Arctic, drying of wetlands may alter the habitat of many species, and may also impact the oil and gas  
24 industry by reducing the availability of water for snow travel and ice roads. In the interior, hydrologic changes  
25 are likely to be felt most acutely in terms of the impacts of drying on fire cycles, as described below, as well as  
26 on changing drainage patterns as permafrost thaws. In western coastal regions, the greatest risks from  
27 changing hydrologic conditions are likely to be erosion (due in part to loss of sea ice and permafrost) and  
28 storms, although current models do not provide clear data on potential changes in storm frequency and  
29 severity. In Southeast Alaska, glacial melt is already occurring, and is likely to continue (citation). This change  
30 in conjunction with changes in the timing of spring runoff, may affect a wide range of ecosystem functions and  
31 human activities, including tourism, hydropower generation, and subsistence harvest.

Comment [FGS1]: Jackie  
Deleted:  
Deleted:  
Deleted: instance  
Deleted: 40  
Deleted: composed of runoff from current rates of systematic ice loss.  
Deleted: 3  
Deleted: 3.0  
Deleted: citation)  
Deleted: .  
Formatted: Font: Not Bold  
Formatted: Font: Not Italic  
Formatted: Font: Not Italic  
Formatted: Font: Not Italic  
Formatted: Font: 9 pt  
Formatted: Font: 9 pt  
Field Code Changed  
Formatted: Font: 9 pt

### 32 Sea Level Change

33 Generally speaking, global sea levels are rising via thermal expansion resulting from warming of the ocean, as  
34 well as freshwater input from the melting of a majority of non-tidewater glaciers and ice sheets in high latitude  
35 zones. In the Gulf of Alaska, for example, approximately 47% of freshwater input is derived from glacier  
36 thinning and ice fields (Neal et al., in review). Consequently, over the next century, global sea levels are  
37 projected to generally rise between 0.6 ft to 1.9 feet (IPCC 2007). However, this projection does not include all  
38 changes in ice-sheet flow or the potential for rapid acceleration in ice loss (Alley et al. 2005, Gregory and  
39 Huybrechts 2006, and Hansen 2005). Several recent studies have found that IPCC estimates of potential sea  
40 level rise might be underestimated regarding ice loss from the Greenland and Antarctic ice sheets (Shepherd  
41 and Wignham 2007, Csatho et al. 2008) and ice loss from mountain glaciers (Meier et al. 2007). Further,

Formatted: Font: 9 pt  
Formatted: Font: 9 pt

1 IPCC results for sea level projections might underestimate sea level rise due to changes in global precipitation  
2 (Wentz et al. 2007, Zhang et al. 2007).

Formatted: Font: Not Bold  
Formatted: Font: Not Italic  
Formatted: Font: Not Italic  
Deleted: however,

3 In some localities of southeast and southcentral Alaska, particularly where glaciers occur in close proximity to  
4 the coastal margin, the land surface is actually rising as a result of the retreat and loss of glacial ice (isostatic  
5 rebound) and secondarily due to active tectonic deformation. This rate of uplift is greater than the projected  
6 rate of global sea level rise and thus, over the next century, the relative sea level in these areas will *decrease*  
7 between 2.1 and 3.4 feet (Larsen et al. 2004, Kelly et al. 2007, Pyare 2009). In contrast, communities in low-  
8 lying areas such as the Yukon-Kuskokwim Delta are likely to face increased flooding and changing storm  
9 surge and storm tracks. In northwest Alaska, overall, decreasing sea ice extent and resulting increasing wave  
10 surge will have greater impact on coastal erosion than sea level rise.

Deleted: 1.0 and 3.6  
Deleted: citation).

11 Ocean Acidification

12 Ocean acidification, occurring as a result of increasing concentrations of CO<sub>2</sub> in the atmosphere, is considered  
13 to be the greatest threat to living marine resources and those who rely on them. The world's largest bodies of  
14 water are effective absorbers of both CO<sub>2</sub> and excess heat associated with climate change, (Sabine et al.  
15 2004). While this has a mitigating effect on the changes to climate, absorption of CO<sub>2</sub> is making the oceans  
16 more acidic posing grave danger to all marine life with calcifying shells. This includes even marine plants and  
17 zooplankton, the basis of the food chain (Feely 2006).

Formatted: Subscript  
Deleted: atmospheric greenhouse gases,  
Deleted: have  
Formatted: Subscript  
Deleted: been effective in absorbing an estimated 30% of global carbon emissions and 80% of the heat generated by the increased levels of greenhouse gases  
Deleted: 's

18 Acidification processes are more pronounced in colder waters (Orr et al. 2005). Since over 50% of the fish  
19 consumed in the U.S. comes from the Bering Sea, this has the potential to significantly alter the conditions  
20 which support commercially important species. Even though it has been noted that some stocks are migrating  
21 further north and into Arctic waters due to ocean warming, acidification is projected to be a challenge there as  
22 well (Orr et al. 2005).

23 Estimates of future atmospheric and oceanic carbon dioxide concentrations suggest that by the middle of this  
24 century atmospheric carbon dioxide levels could reach more than 500 parts-per-million (ppm), and near the  
25 end of the century they could be over 800 ppm. This would result in an additional surface water pH decrease  
26 of approximately 0.3 pH units by 2100 (PMEL n.d.). Box 2-1 describes the effects of ocean acidification on  
27 marine ecosystems.

28

29

Formatted: Font: 9 pt  
Formatted: Font: 9 pt  
Field Code Changed  
Formatted: Font: 9 pt

Formatted: Font: 9 pt

Formatted: Font: 9 pt

### Box 2-1 Ocean Acidification and Its Effect on Marine Species

Deleted: 2

The oceans have absorbed about 50% of the carbon dioxide (CO<sub>2</sub>) released from the burning of fossil fuels, resulting in chemical reactions that lower ocean pH. This has caused an increase in hydrogen ion (acidity) of about 30% since the start of the industrial age through a process known as "ocean acidification." A growing number of studies have demonstrated adverse impacts on marine organisms, including:

- The rate at which reef-building corals produce their skeletons decreases
- The ability of marine algae and free-swimming zooplankton to maintain protective shells is reduced
- The survival of larval marine species, including commercial fish and shellfish, is reduced

This research shows that pH will affect the processes by which animals such as corals, mollusks and crustaceans make their support structures. Because these organisms depend on calcium carbonate, increasing acidity threatens their survival. Pteropods, small planktonic mollusks critically situated at the bottom of the food chain and a key food source for salmon and other species, are particularly vulnerable to increasing acidity in Alaska's marine waters. Other marine organisms at risk from increasing acidification include corals and coralline algae commonly found in reef communities. Cold water coral communities along the Aleutian Islands form important fish habitat. Foraminifera and coccolithophorids, planktons abundant in most surface waters, are also at risk. Some commercial species like clams and crabs will be directly impacted by reductions in calcium carbonate. Others, like most fish populations, will be affected indirectly as acidification impacts their key prey species.

*Continued in next text box...*

1

2

DRAFT

Formatted: Font: 9 pt

Formatted: Font: 9 pt

Field Code Changed

Formatted: Font: 9 pt

Formatted: Font: 9 pt  
Formatted: Font: 9 pt

### Box 2-1 Ocean Acidification and Its Effect on Marine Species (continued)

#### *Ocean Acidification Will Impact Polar and Sub-polar Regions First*

Ocean acidification is likely to alter the biodiversity of the world's marine ecosystems and may affect the total productivity of the oceans. New findings indicate that an increasingly acidic environment could cause problems in high-latitude marine ecosystems within just a few decades. Currently, the oceans' surface water layers have sufficient amounts of calcium carbonate for organisms to use (known as saturated conditions). This calcium carbonate rich layer is deeper in warmer regions and closer to the surface in colder regions. Because calcium carbonate is less stable in colder waters, marine life in the polar oceans will be affected by calcium carbonate loss first. A study published in Nature by 27 U.S. and international scientists stated, "Some polar and sub-polar waters will become under-saturated [at twice the pre-industrial level of CO<sub>2</sub>, 560 ppm], probably within the next 50 years" (Orr et al. 2005).

#### *What Are the Potential Socio-Economic Consequences of Ocean Acidification?*

Ocean acidification will have long-term implications for the global carbon cycle and climate, although the range and magnitude of biogeochemical and biological effects and their socio-economic impacts are currently too uncertain to accurately quantify. However, we do know that such impacts are likely to be substantial.



#### *Feedback to Climate Changes*

As the oceans absorb more and more CO<sub>2</sub> from the atmosphere, their capacity to continue to sequester greenhouse gases diminishes, thereby limiting their ability to reduce the greenhouse effect. This, then inhibits their potential to reduce subsequent warming.



Sources (including photos and chart): PMEL 2008 and Alaska Marine Conservation Council n.d.

1

### 2 Projections – What the Models Predict

3 Scientists extensively use mathematical models of Earth's climate to examine hypotheses about past and  
4 present day climates. These climate simulations provide a framework within which enhanced understanding of

Formatted: Font: 9 pt  
Formatted: Font: 9 pt  
Field Code Changed  
Formatted: Font: 9 pt

Formatted: Font: 9 pt  
Formatted: Font: 9 pt

1 climate-relevant processes, along with improved observations, are merged into coherent projections of future  
2 climate change (CCSP 2008). Over time, these models have evolved into modern coupled atmosphere-ocean  
3 general circulation models (AOGCMs), which incorporate detailed representations of the atmosphere, land  
4 surface, oceans, and sea ice (see Figure 2.3). These models form the basis of the projections for temperature,  
5 precipitation, and other variables reported in the Intergovernmental Panel on Climate Change (IPCC) Fourth  
6 Assessment (IPCC 2007).

Deleted: CITATION

7 Predicting the site-specific magnitude of these changes is crucial to interpreting their impacts. In order to  
8 produce projections that are useful for impacts analysis at the regional level, the coarse-resolution global  
9 climate model output must be "downscaled" to provide higher resolution results. Different methodologies are  
10 available for downscaling, with different degrees of success and reliability (CCSP 2008).

11 For Alaska, the Figure 2-3. Historical Development of Climate Models

Deleted: T

12 Scenarios Network for  
13 Alaska Planning (SNAP)

14 has taken one approach  
15 to developing fine-scale

16 projections of future  
17 climate. SNAP

18 projections are based on  
19 downscaled global

20 models used by the  
21 IPCC). The IPCC used

22 fifteen different GCMs to  
23 prepare its Fourth


24 Assessment Report  
25 (IPCC 2007). Each

26 model was created by a  
27 different nation or group

28 using slightly different  
29 data and assumptions.

30 Thus, models can be  
31 expected to perform with  
32 varying degrees of accuracy in any particular region.

Development of Climate Models: Past, Present, and Future



Source: CCSP 2008.

Deleted: provides

Deleted: in Alaska

Deleted: (citation)

33 To downscale IPCC results to Alaska, SNAP investigators compared model output for past years to actual  
34 climate data for the same time period, and analyzed how well each model predicted monthly mean values for  
35 three different climate variables (surface air temperature, precipitation, and sea level air pressure) over four  
36 overlapping northern regions (Alaska, Greenland, latitude 60-90°N and latitude 20-90°N) for the period from  
37 1958-2000. (Walsh et al. 2008). They noted that models that performed well in one northern region tended to  
38 also perform well in others. SNAP climate models rely on output from the five models that provided the most  
39 accurate overall results. Results are scaled down to match local conditions using data from Alaskan weather  
40 stations and various analytical tools.

Formatted: Font: 9 pt

Formatted: Font: 9 pt

Field Code Changed

Formatted: Font: 9 pt

Formatted: Font: 9 pt  
Formatted: Font: 9 pt

1 Results of SNAP modeling efforts are incorporated into many of the changes to human and natural systems  
2 described below, including vegetation and wildlife species shifts, changes in fire cycles, and loss of  
3 permafrost. Data can be accessed via the SNAP website ([www.snap.uaf.edu](http://www.snap.uaf.edu))

4 Additional linked models connecting climate data to variables such as transportation and construction  
5 parameters, hydrologic shifts, or optimal conditions for tourism, recreation, hunting, and fishing will improve  
6 the connections between SNAP climate data and landscape changes of concern to Alaskans. (Walsh et al.  
7 2008).

### 8 Impacts on Human and Natural Systems in Alaska

9 Alaska is already experiencing climate warming, with key effects occurring to permafrost and sea ice, forests  
10 and other vegetation, coastline communities and infrastructure, marine ecosystems and fisheries, and  
11 subsistence livelihoods (NAST 2000). Over the past three decades, Alaska has experienced a sharp reduction  
12 in snow-cover extent and duration, shorter river- and lake-ice seasons, melting of mountain glaciers, sea-ice  
13 retreat and thinning, permafrost retreat, and increased depth of summer thaw (Weller 2005). The climatic and  
14 other changes associated with warming will not only affect the environment, but will also affect the economy  
15 and daily life and culture in Alaska (see Box 2-2).


#### 16 Permafrost Thawing and 17 Sea Ice Melting

18 Permafrost underlies most of Alaska. Air temperature, snow cover, and vegetation affect the temperature of  
19 the frozen ground and the depth of seasonal thawing. Recent decades of warmer temperatures have  
20 produced extensive warming and thawing of permafrost (Brown and Romanovsky 2008), as illustrated in

21 Figure 2.4. This warming and thawing has  
22 resulted in increased coastal erosion,  
23 landslides, and sinking of the ground surface,  
24 and consequent disruption and damage to  
25 forests, buildings, infrastructure, and coastal  
26 communities (NAST 2000; Nelson et al.  
27 2003). Land subsidence associated with the  
28 thawing of permafrost presents substantial  
29 challenges to engineers attempting to  
30 preserve infrastructure in Alaska (Nelson et al.  
31 2003; Karl et al. 2009, [Larsen et al. 2008](#)).

32 Public infrastructure at risk of damage  
33 includes roads, runways, and water and sewer  
34 systems (Karl et al. 2009). While most  
35 permafrost-related damage to infrastructure is  
36 due to the heat of the infrastructure itself,  
37 most mitigation measures rely on sufficiently  
38 cold winter temperatures to balance the  
39 heating. Many industrial activities (such as oil and gas exploration which requires travel on the tundra) depend  
40 on frozen ground surfaces, and many northern communities rely on ice roads for transport of groceries and

Figure 2-4. Permafrost Temperature, 2978 to 2008.



Source: Karl et al. 2009

Deleted: )

Formatted: Font: 9 pt

Formatted: Font: 9 pt

Field Code Changed

Formatted: Font: 9 pt

Formatted: Font: 9 pt  
Formatted: Font: 9 pt

### Box 2-2. Impacts on the Economy and on People's Lives

The effects of climate change on the environment in Alaska will have consequences for natural resources and production that may affect the Alaskan economy and the lives and livelihood of indigenous and non-indigenous populations.

#### *Impacts on the Economy*

Climate change may affect the economy via effects on natural resources and on transportation and other infrastructure. Large oil and gas reserves exist in Alaska along the Beaufort Sea coast in the Mackenzie River/Beaufort Sea area of Canada. While climate change impacts on this industry have been minor thus far, in the future both negative and positive effects are likely. For example, offshore oil exploration and production is likely to benefit from less extensive and thinner sea ice (Weller 2005). Conversely, ice roads, now used widely for access to offshore activities and facilities, are likely to be less safe and usable for shorter periods (Weller 2005). The thawing of permafrost, on which buildings, pipelines, airfields, and coastal installations supporting oil development are located, is very likely to affect these structures adversely and greatly increase the cost of maintaining or replacing them. (Weller 2005). One study (Larsen et al, 2007, 2008) suggests that strategic adaptation to different future scenarios of climate change could add \$3.6–\$6.1 billion (+10% to +20% above normal wear and tear) to future costs for public infrastructure from now to 2030, and \$5.6–\$7.6 billion (+10% to +12%) from now to 2080.

It is difficult to project impacts on the lucrative Bering Sea fisheries because many factors other than climate are involved (Weller 2005). However, large northward changes in the distribution of fish and shellfish are likely with a warmer climate, which might result in substantial costs to relocate infrastructure supporting commercial fishing (Weller 2005). Warmer waters are likely also to lead to increase primary production in some regions, but a decline in cold-water species such as salmon and pollock.

Other economic sectors in this region, including forestry and agriculture are less developed and less important economically than oil and gas and fish and wildlife. Impacts on these other economic sectors are difficult to assess. For example, impacts on tourism will depend on how the Alaskan features that draw tourists change; large undeveloped landscapes will not be directly affected by climate change, whereas marine mammal populations and accessible glaciers are likely to experience major changes. Impacts on agriculture are also difficult to predict. Growing-degree days have increased by 20%, with benefits for agriculture and forest productivity on some sites, and reduced growth on others (Weller 2005, ACIA 2004).. Moreover, the benefits of a longer growing season will be balanced against the negative effects of decreased soil moisture (Karl et al. 2009).

#### *Impacts on People's Lives*

Traditional lifestyles are already being threatened by the environmental effects associated with climate change, including reduced or displaced populations of marine mammals, seabirds, and other wildlife, and reductions in the extent and thickness of sea ice, making hunting more difficult and dangerous (Weller 2005). Climate change is likely to have significant impacts on the availability of key marine and terrestrial species used as food sources, by shifting the range and abundance of species such as salmon, herring, char, cod, walrus, seals, whales, caribou, moose, and various species of seabird (Weller 2005). Such changes will require major local adjustments in harvest strategies (Weller 2005). These activities play important roles in the lives of many indigenous populations, including making significant contributions to diet and nutrition, providing opportunities for physical activity, contributing monetary income, and providing other important social and cultural functions (Weller 2005).

Impacts have already been observed on lifestyles in indigenous villages and communities in Alaska and Canada that depend heavily on fishing and hunting. Such impacts included reduced access to tundra and offshore food resources, decreases in the anadromous fish stocks and marine mammals harvested for food; threats to villages resulting from coastal erosion, and infrastructure affected by thawing permafrost. Other impacts are likely to occur in the future, due to decreasing area of pack ice, further changes in habitat and migration routes, zoonotic diseases posing a greater threat to humans and wildlife, lower water levels, and other changes.

Formatted: Font: 9 pt  
Formatted: Font: 9 pt  
Field Code Changed  
Formatted: Font: 9 pt

1 other materials. Continued warming will further impair transport by shortening the seasonal use of ice roads

- 1 (ACIA 2004). Thawing is projected to accelerate under future warming, with as much as the top 10 to 30 feet  
 2 of discontinuous permafrost thawing by 2100 (NAST 2000, Romanovsky et al. 2007; Marchenko et al. 2008,  
 3 Romanovsky 2009).
- 4 All components of the cryosphere in the Arctic are experiencing change, including snow cover, mountain and  
 5 continental glaciers, permafrost, sea ice, and lake and river ice (Parson et al. 2000). For example, glaciers in  
 6 Alaska, as throughout the Arctic, have retreated through most of the 20<sup>th</sup> century. Estimated losses in Alaskan  
 7 glaciers are of the order of 30 feet in thickness over the past 40 years (Parson et al. 2000).

Figure 2-5. Sea Ice Extent in 2008 and



Source: Richter-Menge et al. 2008

Sea ice off the Alaskan Coast is retreating and thinning, with widespread effects on marine ecosystems, coastal climate, human settlements, and subsistence activities. Recent studies estimate arctic-wide reductions in annual average sea-ice extent of about 5-10% and a reduction in average thickness of about 10-15% over the past few decades (ACIA 2004). Figure 2-5 depicts the extent of sea ice in September 2008. The magenta line represents median sea ice extent between 1979 and 2008. The annual maximum sea ice extent typically occurs in March. This annual maximum level has been gradually declining since 1979, and experienced a record low in 2006, with a slight rebound since the low (Richter-Menge et al. 2008). Retreat of sea ice allows larger storm surges to develop, increasing the risk of inundation and increasing erosion on costs already made vulnerable by permafrost thawing (NAST 2000).

24

- 25 Loss of sea ice also causes large scale changes in marine ecosystems, and threatens populations of marine  
 26 mammals and polar bears that depend on ice (NAST 2000).
- 27 At the same time, the continued reduction of sea ice is very likely to increase the navigation season, and

Figure 2-6. Coastline Erosion in Shishmaref, Alaska.



Shishmaref, where the coastline has eroded 100-300 feet in the past 30 years. Source: The Nome Nugget

seasonal opening of the Northern Sea Route is likely to make trans-arctic shipping feasible during summer months, within several decades, although increasing ice movement will initially make shipping more difficult in some channels of the Northwest Passage.

Lakes are also declining in area. Across the southern two-thirds of Alaska, the area of closed-basin lakes (lakes without stream inputs and outputs) has decreased over the past 50 years (Karl et al. 2009). This is likely due to the greater evaporation and thawing of permafrost that result from warming (Karl et al. 2009). Continued decline in the area of surface water would present challenges for the management of natural

Formatted: Font: 9 pt  
Formatted: Font: 9 pt

### Box 2-3 Alaska Native Villages: Limited Progress Has Been Made on Relocating Villages Threatened by Flooding and Erosion

In December 2003, GAO reported that most of Alaska's more than 200 Native villages were affected to some degree by flooding and erosion (GAO 2003). Since 2003, state officials have identified the growing impacts of climate change, increasing the urgency of federal and state efforts to identify imminently threatened villages and assess their relocation options.

Since 2003, federal, state, and village officials have identified 31 villages that face imminent threats. The U.S. Army Corps of Engineers' (Corps) March 2009 Alaska Baseline Erosion Assessment identified many villages threatened by erosion, but did not assess flooding impacts. At least 12 of the 31 threatened villages have decided to relocate—in part or entirely—or to explore relocation options.

Federal programs to assist threatened villages prepare for and recover from disasters and to protect and relocate them are limited and unavailable to some villages. In the absence of a lead entity, federal agencies individually prioritize assistance to villages on the basis of their programs' criteria. These criteria do not necessarily ensure that the villages in greatest peril get the highest priority, and although the Corps has assessed erosion threats, there is no lead federal entity to prioritize and coordinate assistance using this information.

Of the 12 villages exploring relocation options, Newtok has made the most progress in its relocation efforts. The Newtok Planning Group, formed in 2006 by federal, state, regional, and village partners, has helped to accelerate the relocation process that the village proactively initiated in 1994. In 2007, the Newtok Planning Group reported that the lack of designated federal and state lead entities to guide, coordinate, and fund assistance impeded village relocation efforts and created uncertainty regarding the fulfillment of environmental analysis requirements under the National Environmental Policy Act. In 2008, the state designated a lead agency for village relocation assistance, and federal, state, and village officials told GAO that a similar lead federal entity is needed. Lead authority could be provided to an existing agency or commission, or a new entity could be formed for this purpose.

Source: GAO 2009

1 resources and ecosystems on National Wildlife Refuges in Alaska (Karl et al. 2009). These refuges, which  
2 cover 77 million acres (21 percent of Alaska) and comprise 81 percent of the US national Wildlife Refuge  
3 System, provide breeding habitat for millions of waterfowl and shorebirds that winter in the lower 48 state (Karl  
4 et al. 2009).

#### 5 Threats to Coastal Communities, 6 Habitats, and Fishing Fleets

7 Alaska has more coastline than the other 49 states combined (Karl et al. 2009). Coastal erosion is causing the  
8 shorelines of some areas to retreat at average rates of tens of feet per year (Karl et al. 2009). Frequent storms  
9 in the Gulf of Alaska and the Bering, Chukchi, and Beaufort Seas already affect the coasts during much of the  
10 year (Karl et al. 2009). Alaska' coastlines, many of which are low in elevation, are increasingly threatened by a  
11 combination of the loss of their protective sea ice buffer, increasing storm activity, and thawing coastal  
12 permafrost (Karl et al. 2009). Increases in the frequency and intensity of storm surges have triggered  
13 increased coastal erosion that is threatening a number of coastal villages (see Figure 2-6). A recent report  
14 from the General Accountability Office (GAO) indicated that, since 2003, federal, state, and village officials  
15 have identified 31 villages that face imminent threats (see Box 2-3).

16 Storm surges have also reduced the protection that barrier islands and spits provide to coastal habitats.  
17 Commercial fishing fleets and other marine traffic are also strongly affected by Bering Sea storms. High-wind

Formatted: Font: 9 pt  
Formatted: Font: 9 pt  
Field Code Changed  
Formatted: Font: 9 pt

1 events have become more frequent along the western and northern coasts (Karl et al. 2009). The same  
2 regions are experiencing increasingly long sea-ice-free seasons and hence longer periods during which  
3 coastal areas are especially vulnerable to wind and wave damage (Karl et al. 2009).

#### 4 Forest and Vegetation Changes

5 The Arctic region, particularly Alaska, is already experiencing major ecological impacts of warming. Rising  
6 temperatures have caused northward expansion of boreal forest in some areas, significant increases in fire  
7 frequency and intensity, and unprecedented insect outbreaks (Weller 2005, ACIA 2004). During the 1990s, for  
8 example, south-central Alaska experienced the largest outbreak of spruce beetles in the world, attributable to  
9 the combination of rising temperatures speeding up the life cycle of the beetle and extended drought  
10 weakening the trees (Karl et al. 2009). Large areas of dead trees, such as those left behind by pest  
11 infestations, are highly flammable and more vulnerable to wildfire than living trees (Karl et al. 2009).

12 Climate plays a key role in determining the extent and severity of wildfires, as well as insect outbreaks (Karl et  
13 al. 2009). The amount of boreal forest in North America that burned annually tripled from the 1960s to the  
14 1990s, and half of the severe fire years on record have occurred since 1990 (Kasischke and Turetsky 2006).  
15 Fire in Alaska is tightly linked to climate (Duffy 2005) and the average area burned per year in Alaska is  
16 projected to double by the middle of this century (Balshi et al. 2008). Under a moderate projection of climate  
17 change, by the end of this century, the total area burned by fire is projected to triple (Karl et al. 2009). Current  
18 projections suggest that, due to increases in burn area per decade, the tundra-dominated landscape on  
19 Seward Peninsula will eventually be replaced by deciduous forest (ACIA 2004). In other areas, forested areas  
20 are likely to convert to bogs as permafrost thaws (ACIA 2004). Coupled trajectories of future climate and  
21 demographics, direct, near-term threats to life and property from wildfire in Alaska are expected to increase  
22 (Trainor et al 2009).

#### 23 Sensitivity of Marine 24 Ecosystems and Fisheries

25 The Gulf of Alaska and Bering Sea support marine ecosystems of great diversity and productivity. Alaska  
26 leads the United States in the value of its commercial fishing catch, and most of the nation's salmon, crab,  
27 halibut, and herring come from Alaska (Karl et al. 2009). One of the most productive areas for Alaska fisheries  
28 is the northern Bering Sea off Alaska's west coast (Karl et al. 2009). Recent climate-related impacts observed  
29 in the Bering Sea include significant reductions in seabird and marine mammal populations, unusual algal  
30 blooms, abnormally high water temperatures, and low harvests of salmon on their return to spawning areas  
31 (Weller 2005). While the Bering Sea Fishery has become one of the world's largest, the area has been  
32 undergoing change for several decades. The abundance of Stellar sea lions has declined by between 50%  
33 and 80%, and northern fur seal pups on the Pribilof Islands – the major Bering Sea breeding grounds –  
34 declined by 50% between the 1950s and 1980s (Weller 2005, ACIA 2004). Populations of some seabird  
35 species, including common the murre, have also declined significantly (Weller 2005).

36 Differentiating among the various factors affecting the Bering Sea ecosystem is a major focus of current and  
37 projected research (Weller 2005). However, future projections for the Bering Sea suggest productivity  
38 increases at the base of the food chain, poleward shifts of some cold-water species, and negative effects on  
39 ice-dwelling species (Weller 2005, ACIA 2004). Warmer temperatures will also affect commercial fisheries,  
40 with large northward shifts of fish and shellfish species, and associated declines in production of cold-water  
41 species such as salmon and pollock, and increased production of other species (ACIA 2004).

Formatted: Font: 9 pt  
Formatted: Font: 9 pt

1 Changes In the Diversity, Ranges,  
2 and Distributions of Species

3 The Arctic sub-region that includes Alaska, Chukotka, and the Western Canadian Arctic, is home to over 70  
4 percent of the rare plant species that occur only in the Arctic, and home to a number of plant and animal  
5 species already classified as "threatened" (ACIA 2004). Species concentrated in small areas, such as  
6 Wrangell Island, are particularly  
7 vulnerable to the direct effects of climate  
8 change, combined with competition from  
9 migrating non-native species (ACIA  
10 2004).

11 Increased Stress on  
12 Subsistence  
13 Livelihoods and Lifestyles

14 Subsistence makes an important  
15 contribution to livelihood in many  
16 isolated rural communities, especially  
17 but not exclusively for native peoples.  
18 Livelihoods that sustain indigenous  
19 communities include hunting, trapping,  
20 gathering, and fishing. These activities  
21 not only make significant contributions  
22 to the diet and health of many  
23 indigenous populations, but also play  
24 large and important social and cultural  
25 roles (ACIA 2004). Reduced or  
26 displaced populations of marine  
27 mammals, seabirds, and other wildlife,  
28 together with continuing thinning of sea-  
29 ice, have affected the safety, and the  
30 dietary and economic well-being of  
31 subsistence communities (ACIA 2004).  
32 Over the course of this century, the  
33 most productive commercial fisheries  
34 are likely to become more distant from  
35 existing fishing ports and processing  
36 infrastructure (Karl et al. 2009). These  
37 changes will also affect the ability of  
38 Native Peoples to successfully hunt and fish for food. Particularly for rural communities, adapting to climate  
39 change will pose some complex challenges (see Box 2-5).

**Box 2-4. Vulnerability to Climate Change and the Arctic**

The majority of the Arctic's residents live in small to medium-sized communities, in many cases located in remote regions and dependent on climate-sensitive livelihoods, including hunting, fishing, herding and forestry. This dependence on climate-sensitive resources and infrastructure will make the Arctic particularly sensitive to climate change. Climate change is already threatening activities such as hunting, and raising questions about the long-term sustainability of traditional way.

In the Arctic, adaptation is increasingly prominent in policy discussions, with national and regional governments, non-governmental organizations, communities, and national and international research bodies stressing the need to strengthen the ability of communities, regions, and economic sectors to adapt to current and future climate change.

To identify adaptation needs and inform the development of policies to reduce the negative impacts of climate change, it is crucial to identify and characterize vulnerability. Vulnerability can be thought of as the capacity to be wounded: it is a measure of the susceptibility to harm in a system in response to a stimulus or stimuli. In this context, the stimuli are climate-related risks, and the "system" can range from an individual or household unit to the national state. In turn, vulnerability is related to both exposure and sensitivity to climatic risks, and the adaptive capacity to deal with those risks.

Non-climatic factors can amplify or attenuate vulnerability to climatic stress. These include sources of livelihoods, assets, access to resources, globalization, institutional networks, education, gender, race, ethnicity, and poverty. These determinants are influenced by social, economic, cultural and political conditions. Identifying viable options for adaptation requires both identifying technological and engineering-based responses and also understanding and applying the lessons of vulnerability science, to the Arctic.

Source: Ford and Furgal 2009.

Deleted: term

Formatted: Font: 9 pt  
Formatted: Font: 9 pt  
Field Code Changed  
Formatted: Font: 9 pt

## 1 **Technical Working Groups Supporting Alaska's Adaptation Advisory** 2 **Group**

3 Prior to establishing Technical Work Groups (TWG) to help the AAG formulate options, the Sub-Cabinet  
4 conducted an inventory of the potential ecological and socioeconomic impacts of climate change on Alaska. A  
5 thorough analysis of peer-reviewed literature, such as the Arctic Climate Impact Assessment (ACIA 2005),  
6 was conducted along with a review of impacts identified by appropriate state agencies, and the compilation  
7 put together by the Alaska Climate Impact Assessment Commission (ACIAC 2008). This laid the groundwork  
8 for categorizing the impacts into sectors that formed focal points for the Technical Work Groups. The AAG  
9 formed the TWGs to develop options to address potential impacts in four sectors: Health and Culture, Public  
10 Infrastructure, Natural Systems and Other Economic Activities. The four sectors are not entirely discrete; over  
11 the course of the strategy development process, the AAG and TWGs dealt with a number of issues in the  
12 scope and potential overlaps between the impact categories.<sup>1</sup>

### 13 **Health and Culture**

14 This sector focuses on the human health, cultural, recreational, and quality-of-life impacts of climate change  
15 on Alaskans. Climate change is projected to be associated with increases in some diseases that flourish in  
16 warmer temperatures (such as paralytic shellfish poisoning). Water quality and availability—a decrease in  
17 quality of potable water due to drought, saltwater intrusion, and other issues—will also have health  
18 consequences. The health and culture of subsistence lifestyles are especially at risk, due to diminishment or  
19 change of the subsistence diet, due, in part, to reduced availability of traditional food supplies (waterfowl, fish,  
20 marine mammals, sea vegetables, berries, and plant medicines). Injuries, due to thinning and other changes  
21 in ice, wildfires, and insect bites and stings, are also likely to increase.

### 22 **Natural Systems and Associated Economies**

23 This sector examines the impacts of climate change on biodiversity, ecosystem health, and associated human  
24 economic activity. In general, climate is projected to result in changes in growing seasons, and changes in the  
25 distribution, quantity, and ranges of many species. Consequently, this sector comprises a wide range of  
26 subsectors, including agriculture, boreal and temperate forests and dependent species, forestry, tundra and  
27 alpine ecosystems and dependent species, freshwater ecosystems and dependent species, marine, sea ice,  
28 coastal environment, and dependent species, other warm temperature impacts on animals, commercial and  
29 sport fishing, subsistence, fishing, hunting, trapping, and gathering, sport hunting, tourism and watchable  
30 wildlife.

### 31 **Public Infrastructure**

32 This sector addresses the physical impacts of climate change on Alaska's built environment and  
33 transportation options. Infrastructure includes both publicly- and privately-owned infrastructure (the word  
34 public refers to "use" rather than "ownership"). Types of infrastructure considered by this sector include road  
35 transportation (highways, roads, and bridges), air transportation (airports and landing strips), sea walls and  
36 shoreline protection, utility and fuel infrastructure, landfills, sewage and septic systems, and water systems.

---

<sup>1</sup> Additional information on the initial characterization of sectors can be found on Alaska's Climate change website, associated with the first meeting of the AAG. <http://www.climatechange.alaska.gov/aag/aag.htm>

1 As discussed above, impacts from thawing permafrost, increased freeze-thaw cycles, and erosion all pose  
 2 challenges to road infrastructure and buildings. Other forms of infrastructure are similarly susceptible to these  
 3 and other changes, including increased fire risk, increased coastal fog (affecting flying conditions), and other  
 4 impacts.

#### 5 Other Economic Activities

6 This sector focuses on Alaskan economies affected by a changing climate that are not directly dependent on  
 7 living ecosystems. Key industries initially examined by the TWG included oil and gas, mining, ocean  
 8 transportation and other transportation, tourism, and other business impacts and opportunities. The types of  
 9 impacts addressed by this sector include the effects of climate change on industries such as oil and gas,  
 10 which are expected to be negatively affected by difficulties in tundra ice travel and effects of sea level rise and  
 11 erosion on buried or above-ground oil and gas pipelines. The sector also focuses on economic opportunities,  
 12 such as increased shipping opportunities.

13 Figure 2-7 provides an overview of the four sectors and the issues covered by each TWG.

14

Figure 2-7. Effects of Climate Change on Each Sector.

